

Is Schrödinger's Conjecture for the Hydrogen Atom Coherent States Attainable?

Ivaylo Zlatev¹, Wei-Min Zhang² and Da Hsuan Feng^{1,3}

¹*Department of Physics, Drexel University, Philadelphia, PA 19104-9984 USA*

²*Institute of Physics, Academia Sinica, Taipei 11529, Taiwan, ROC*

³*Physics Division, Oak Ridge National Laboratory, Oak Ridge, TN 37831 USA*

February 9, 2008

Abstract

We construct the most general $SO(4,2)$ hydrogen atom coherent states which are the counterpart of Schrödinger's harmonic oscillator coherent states. We show that these states cannot be localized and cannot follow the classical orbits. Thus, Schrödinger's conjecture for the hydrogen atom coherent states is unattainable.

The classical limit of any quantum system is an important open question. A natural way to realize the quantum to classical transition, first proposed by Schrödinger in 1926[1], is via coherent states [2]. These states, at least for the harmonic oscillator (HO), are well-defined localized wavepackets that move along the corresponding classical orbits. In the last paragraph of his paper, Schrödinger postulated that perhaps states with similar properties may be constructed for the other simplest yet fundamental quantum system, the hydrogen atom, and that the wavepackets for this system would travel along Kepler elliptic orbits. He pointed out that “the technical difficulties in the calculation (were) greater” than in the HO case. However, sixty-seven years later, such states *have not* yet been constructed even though there were numerous attempts. The main conclusion of this paper is to point out that in principle such states as envisioned by Schrödinger *do not exist*.

The HO coherent states Schrödinger constructed are linear superpositions of all eigenstates of the HO which form the irreducible representation space (Fock space) of the Heisenberg-Weyl group H_4 (the dynamical group of the HO) [2]. All hydrogen atom bound eigenstates (including all the degenerate states) form an irreducible representation space of the $SO(4,2)$ dynamical group [3]. Thus, to find the analogous hydrogen atom coherent states that are linear superpositions of all its eigenstates, it is natural to use the group theoretical approach to construct the $SO(4,2)$ coherent states for the hydrogen atom.

A set of coherent states were previously constructed from the $SO(4,2)$ group and it was claimed that they follow the Kepler motion [4]. The validity of this statement was later questioned [5]. There were of course other attempts to construct the coherent states of the hydrogen atom by the use of wavepackets which did not utilize the above irreducible representation space of $SO(4,2)$ [6, 7, 8]. In this paper we will show that the $SO(4,2)$

coherent states of the hydrogen atom can neither satisfy the general minimum uncertainty relation, nor follow the Kepler orbits. Since these are the most general coherent states for the system, we are compelled to conclude that the hydrogen atom counterpart of Schrödinger's HO coherent states cannot possess the same physical features as the HO states.

The $SO(4,2)$ group is a 3-rank, 15-dimensional non-compact semisimple Lie group. Its generators in the hydrogen atom representation are [3]:

$$\begin{aligned} \mathbf{L} &= \mathbf{r} \times \mathbf{p}, \quad \mathbf{A} = \frac{1}{2}\mathbf{r}p^2 - \mathbf{p}(\mathbf{r} \cdot \mathbf{p}) - \frac{1}{2}\mathbf{r}, \\ \mathbf{M} &= \frac{1}{2}\mathbf{r}p^2 - \mathbf{p}(\mathbf{r} \cdot \mathbf{p}) + \frac{1}{2}\mathbf{r}, \quad \mathbf{\Gamma} = r\mathbf{p}, \\ S &= \frac{1}{2}(rp^2 - r), \quad T = \mathbf{r} \cdot \mathbf{p} - i, \quad \Gamma_0 = \frac{1}{2}(rp^2 + r). \end{aligned} \tag{1}$$

The irreducible representation space for the hydrogen atom is a discrete lower-bound space with basis $|nlm\rangle$ which are the eigenstates of the hydrogen atom. The lowest state is $|\psi_0\rangle \equiv |n=1, l=0, m=0\rangle$. The maximum subgroup of $SO(4,2)$ which leaves the ground state invariant is $SO(4) \otimes SO(2)$. Its generators are L_i , A_i and Γ_0 . According to the general algorithm [2], the $SO(4,2)$ coherent states for the hydrogen atom are then defined as:

$$\begin{aligned} |SO(4,2), CS\rangle &\equiv |\Omega\rangle = \Omega|\psi_0\rangle \\ &= \exp i(\alpha \cdot \mathbf{M} + \beta \cdot \mathbf{\Gamma} + \alpha_4 S + \beta_4 T) |\psi_0\rangle, \end{aligned} \tag{2}$$

where $\{\alpha_i, \beta_i; (i=1, 2, 3, 4)\}$ are real parameters.

These coherent states for the hydrogen atom have useful algebraic and topological properties. Just as the HO coherent states are isomorphic to the coset space $H_4/U(1) \otimes U(1)$, these states are isomorphic to the coset space $SO(4,2)/SO(4) \otimes SO(2)$, a 4-dimensional complex manifold which provides the geometrical basis for the link with

the classical limit of the hydrogen atom. Their cardinal algebraic property is the resolution of identity:

$$\int d\mu |\Omega\rangle\langle\Omega| = I \quad (3)$$

where $d\mu$ is the normalized $SO(4,2)$ -invariant measure on $SO(4,2)/SO(4) \otimes SO(2)$. This property offers a connection between the quantum observables and the classical observables [9]. We mentioned that there were other constructions of wavepackets for the hydrogen atom [6, 7, 8]. These wavepackets do not satisfy the above resolution of identity [9].

One can physically appreciate the coherent states we have constructed for the hydrogen atom by comparing them to the Schrödinger's HO coherent states. It is now well known that the Schrödinger's HO coherent states can be written as [10]:

$$|z\rangle = \exp(za^\dagger - z^*a)|\psi_0\rangle = e^{-|z|^2/2} \sum_{n=0}^{\infty} \frac{z^n}{\sqrt{n!}} |n\rangle \quad (4)$$

where $|n\rangle$ are the HO eigenstates. These coherent states are wavepackets and are well localized at the point z in the HO phase space. This is a direct result of the fact that $\frac{z^n}{\sqrt{n!}}$ has its maximum at $|z|^2 = n$, as Schrödinger pointed out in his paper. Hence, in the expansion, the coherent states will be dominated by a small number of eigenstates in the neighborhood of this $|z|^2$. Here we must point out that this is true not only for very large $|z|$ as Schrödinger had stated in his paper, but for any value of $|z|$ as well. The rigorous statement of the above well-localized wavepackets is that they will satisfy the minimum uncertainty relation:

$$\Delta q \Delta p = \hbar/2. \quad (5)$$

Schrödinger stated in his paper [1] that “*das man auf ganz ähnliche Weise auch die Wellengruppen konstruieren kann, welche auf hoch-quantigen Keplerellipsen umlaufen und das undulationsmechanische Bild des Wasserstoffelektrons sind.*” [11]. The word

“hoch-quantingen” means “high quantum number” [12]. The same sort of statement about the HO case was made earlier by Schrödinger in his paper, and we now see that in fact it is not necessary for the HO coherent states. Hence, we should not expect that the equivalent hydrogen atom coherent states need to be constructed from eigenstates with high quantum numbers.

To compare the above picture of the HO coherent states to that of hydrogen atom coherent states, it is convenient to use the boson representation of the $SO(4,2)$ algebra [3]:

$$\begin{aligned} L_k &= \frac{1}{2}\epsilon_{ijk}(\mathbf{a}^\dagger\sigma_k\mathbf{a} + \mathbf{b}^\dagger\sigma_k\mathbf{b}), \quad A_i = -\frac{1}{2}(\mathbf{a}^\dagger\sigma_i\mathbf{a} - \mathbf{b}^\dagger\sigma_i\mathbf{b}), \\ M_i &= -\frac{1}{2}(\mathbf{a}^\dagger\sigma_i C\mathbf{b}^\dagger - \mathbf{a}C\sigma_i\mathbf{b}), \quad \Gamma_i = -\frac{i}{2}(\mathbf{a}^\dagger\sigma_i C\mathbf{b}^\dagger + \mathbf{a}C\sigma_i\mathbf{b}), \\ S &= \frac{1}{2}(\mathbf{a}^\dagger C\mathbf{b}^\dagger + \mathbf{a}C\mathbf{b}), \quad T = -\frac{i}{2}(\mathbf{a}^\dagger C\mathbf{b}^\dagger - \mathbf{a}C\mathbf{b}), \quad \Gamma_0 = \frac{1}{2}(\mathbf{a}^\dagger\mathbf{a} + \mathbf{b}^\dagger\mathbf{b} + 2), \end{aligned} \quad (6)$$

where σ_i are the Pauli spin matrices, $C = i\sigma_2$ and $\mathbf{a} = (a_1, a_2)$, $\mathbf{b} = (b_1, b_2)$ are the two harmonics oscillator creation and annihilation operators. The hydrogen atom ground state is the boson vacuum state, $a_i|\psi_0\rangle = b_i|\psi_0\rangle = 0$, $i = 1, 2$, and its coherent states, Eq.(2), can then be reexpressed as

$$|\Omega\rangle = \exp \sum_{i,j=1}^3 \left(z_{ij} a_i^\dagger b_j^\dagger - z_{ij}^* a_i b_j \right) |\psi_0\rangle, \quad (7)$$

where the relations between z_{ij} and $\{\alpha_i, \beta_i\}$ can be found by comparing Eq.(2) and Eq.(7).

The coherent states can further be rewritten as

$$\begin{aligned} |\Omega\rangle &= \mathcal{N}^{1/2} \sum_{n=1}^{\infty} \sum_{k_1 k_2 k_3 k_4} C_{1234} \frac{(z_{11}^{k_1} z_{12}^{k_2} z_{21}^{k_3} z_{22}^{k_4})}{(n-1)!} |n\rangle \\ &= \mathcal{N}^{1/2} \sum_{n=1}^{\infty} \sum_{lm} C_{lm}^z |nlm\rangle \end{aligned} \quad (8)$$

where $\mathcal{N}$ is the normalized constant, $k_1 + k_2 + k_3 + k_4 = n - 1$, and $|n\rangle = (a_1^\dagger)^{k_1+k_2} (a_2^\dagger)^{k_3+k_4} (b_1^\dagger)^{k_1+k_3} (b_2^\dagger)^{k_2+k_4} |\psi_0\rangle$ which are the eigenstates of the hydrogen atom

with the principal quantum number n

$$\Gamma_0|n\rangle = n|n\rangle. \quad (9)$$

Using Schrödinger's argument for the HO coherent states, we can see that the $SO(4,2)$ coherent states for hydrogen atom are not well-localized wavepackets. This is because from Eq.(8) it is easy to see that for given z_{ij} , there is no unique peak as in the HO case. On the contrary, there is a large number of peaks which will *de facto* delocalize the wavepackets. Hence it is not expected that the $SO(4,2)$ coherent states for the hydrogen atom can satisfy the minimum uncertainty relation.

Next we will show rigorously that the $SO(4,2)$ coherent states cannot satisfy the general minimum uncertainty:

$$(\Delta r)^2(\Delta p_r)^2 = \frac{1}{4}\langle\Omega|[r - \langle r\rangle, p_r - \langle p_r\rangle]_+|\Omega\rangle^2 + \frac{\hbar^2}{4}, \quad (10)$$

of which the Heisenberg minimum uncertainty relation is a special case. The proof is as follows. Eq.(10) is true only if [13]:

$$(r - \langle r\rangle)|\Omega\rangle = c(p_r - \langle p_r\rangle)|\Omega\rangle, \quad (11)$$

which is the same as:

$$(r^2 - r\langle r\rangle)|\Omega\rangle = c(\mathbf{r} \cdot \mathbf{p} - r\langle p_r\rangle)|\Omega\rangle. \quad (12)$$

However, we find that Eq.(12) cannot be satisfied by the $SO(4,2)$ coherent states. Using the matrix representation approach in the coherent state calculation [2], Eq.(12) can be reduced to

$$A|0\rangle + B|2\rangle + C|4\rangle = 0 \quad (13)$$

where $|0\rangle \equiv |\psi_0\rangle$, $|2\rangle \equiv a_i^\dagger b_j^\dagger |\psi_0\rangle$, and $|4\rangle \equiv a_i^\dagger b_j^\dagger a_k^\dagger b_l^\dagger |\psi_0\rangle$, and the coefficients A, B, C are functions of z_{ij} . It can explicitly be shown that the coefficients A, B and C vanish only

if $C = \frac{\langle r^2 \rangle}{\langle \mathbf{r} \cdot \mathbf{p} \rangle}$, and

$$2Y_1X + iX\sigma_2X - iY_1\sigma_2Y_2 = 0, \quad X\sigma_2X + Y_1\sigma_2Y_2 = 0, \quad (14)$$

where

$$X = \frac{Z}{\sqrt{Z^\dagger Z}} \sinh \sqrt{Z^\dagger Z}, \quad (15)$$

Z is a 2×2 matrix with elements z_{ij} , the parameters in the coherent states of Eq.(7), and

$$Y_1 = \sqrt{I + XX^\dagger}, \quad Y_2 = \sqrt{I + X^\dagger X} \quad (16)$$

Now using Eq.(14), we find that $\Delta r = 0$, and Δp_r is infinite, which seems to be consistent with the uncertainty relation. However, in this case, the coherent state becomes an eigenstate of the r operator with zero eigenvalue and therefore contains no physical information. In fact, we find that Eq.(14) leads to

$$-4Im^2(x_{12}) - 1 = (Re(x_{11}) + Re(x_{22}))^2 + Im(x_{11}) - Im(x_{22}))^2 \quad (17)$$

a condition which cannot be satisfied. In eq. (17), $Re(x_{ij})$ and $Im(x_{ij})$ mean the real and imaginary part of the matrix element x_{ij} of the matrix X . This indicates that Eq.(10) cannot be obeyed. Thus, the requirement that the $SO(4,2)$ coherent states satisfy the general minimum uncertainty cannot be realized.

We conclude that the most general $SO(4,2)$ coherent states for the hydrogen atom cannot be well-localized wavepackets. In fact, based on the recent work proposed by two of the authors [14], we can show that the coherent states that are the superposition of all the eigenstates of the hydrogen atom cannot give us the classical limit. The classical limit in this framework is realized as follows: classical dynamics, if it exists at all for a quantum system, is depicted by the phase space structure of the expectation value of the hamiltonian in the coherent states in which the quenching index Ξ , which is inherently

hidden in the coherent states, goes to infinity. In the $SO(4,2)$ coherent states for the hydrogen atom, $\Xi = \frac{1}{2}$. So, there is no classical limit for this quantum system. As a result, the coherent states cannot move along the Kepler orbits. It would therefore appear that the goal of Schrodinger is unattainable.

Finally, we should remark that the results we have obtained here do not preclude that coherent states formed by a complete *subset* of the hydrogen eigenstates have a classical limit. Such states are for example the $SU(1,1)$ coherent states, which are superpositions of the states with $n \geq l + 1$ where l is the angular momentum quantum number and should be a relatively large value since the quenching index for $SU(1,1)$ hydrogen atom coherent states is $\Xi = 2(l + 1)$ [14]. Quite recently, there were some remarkable experimental developments on single atoms and radiation [5, 15] which may imply that there will be some wavepackets which will follow classical trajectories and have both classical and quantum properties. Our work on the $SU(1,1)$ hydrogen atom coherent states is now in progress. We speculate that these could be the $SU(1,1)$ coherent states for the hydrogen atom.

This work is supported by the U.S. National Science Foundation and the National Science Council of the ROC. We thank B. Giraud for a useful communication.

References

- [1] E. Schrödinger, *Naturwissenschaften*, **4**, 664 (1926).
- [2] For a review, see Wei-Min Zhang, Da Hsuan Feng, and R. Gilmore, *Rev. Mod. Phys.* **62** 867 (1990), and references therein.
- [3] For a complete discussion of the hydrogen atom dynamical group see A. O. Barut and G. Bornzin, *J. Math. Phys.* **12**, 841 (1971), and B.G. Wybourne, *Classical Groups for Physicists*, (New York, Wiley, 1974), p.311.
- [4] J. Mostowski, *Lett. Math. Phys.* **2**, 1,(1977).
- [5] Z.D. Gaeta, C.R. Stroud Jr., *Phys. Rev. A*, **42**, 6308 (1990).
- [6] L.S. Brown, *Am. J. Phys.*, **41**, 525 (1973).
- [7] J. A. Yeazell, C.R. Stroud Jr., *Phys. Rev. A*, **43**, 5153 (1991).
- [8] for example J.C. Gay et al., *Phys. Rev. A*, **39** 6587 (1989) and references therein; M. Nauenberg, *Phys. Rev. A* **40** 1133 (1989).
- [9] Coherent states must satisfy the resolution of identity. See the discussion in the preface of the book *Coherent states* edited by J. R. Klauder and B. S. Skagerstam (World Scientific, Singapore, 1985).
- [10] This coherent state is equivalent to expression (4) in [1]. See R. Glauber, *Phys. Rev.* **130**, 2529 (1963)
- [11] In English, it is “in a similar way, wave groups can be constructed which move round highly quantised Kepler ellipses and are the representation by wave mechanics of the hydrogen electron” see *Collected papers on Wave Mechanics* (Blackie & Son Limited, London, 1958)
- [12] We thank F. K. Thielemann for the precise translation.
- [13] R. Shankar, *Principles of Quantum Mechanics*, (Plenum Press,1980).
- [14] Wei-Min Zhang and Da Hsuan Feng, *J. Mod. Phys. Lett.*, **8**, 1417 (1993).

- [15] see for example J. A. Yeazell, M. Mallalieu, and C.R.Stroud, Jr., Phys. Rev. Lett. **64**, 2007, (1990), L. Moorman, P.M. Koch, in *Quantum Non-Integrability*, edited by D. H. Feng and J. M. Yuan, (World Scientific, Singapore, 1992).